


---

---

# FIELDIANA

---

---



## Geology

NEW SERIES, NO. 52

### A Paleopopulation of *Coryphodon lobatus* (Mammalia: Pantodonta) from Deardorff Hill *Coryphodon* Quarry, Piceance Creek Basin, Colorado

Elizabeth M. McGee

William D. Turnbull



January 8, 2010  
Publication 1554

---

PUBLISHED BY FIELD MUSEUM OF NATURAL HISTORY

---

## Mission

*Fieldiana* is a peer-reviewed monographic series published by the Field Museum of Natural History. *Fieldiana* focuses on mid-length monographs and scientific papers pertaining to collections and research at The Field Museum. The four series pertain to subject matter in the fields of Anthropology, Botany, Geology, and Zoology.

## Eligibility

Field Museum curators, research associates, and full-time scientific professional staff may submit papers for consideration. Edited volumes pertaining to Field Museum collections may also be submitted for consideration under a subsidy arrangement. The submission and peer review of these chaptered volumes should be arranged well in advance with the managing scientific editor and the appropriate associate editor.

## Submission Procedures

Submission procedures are detailed in a separate document called "SUBMISSIONS PROCEDURES" available on the *Fieldiana* web site: ([http://www.fieldmuseum.org/research\\_collections/fieldiana/](http://www.fieldmuseum.org/research_collections/fieldiana/)) under the Author's page. All manuscripts should be submitted to the managing scientific editor.

## Editorial Contributors:

### Managing Scientific Editor

Harold K. Voris ([hvoris@fieldmuseum.org](mailto:hvoris@fieldmuseum.org))

### Acting Editorial Coordinator

Peter Lowther ([plowther@fieldmuseum.org](mailto:plowther@fieldmuseum.org))

### Editorial Assistant

Chris Jones ([cjones2@fieldmuseum.org](mailto:cjones2@fieldmuseum.org))

### Illustration Editor

Lisa Kanellos ([lkanellos@fieldmuseum.org](mailto:lkanellos@fieldmuseum.org))

## Anthropology



### Co-Associate Editors

Jonathan Haas ([jhaas@fieldmuseum.org](mailto:jhaas@fieldmuseum.org))

Gary Feinman ([gfeinman@fieldmuseum.org](mailto:gfeinman@fieldmuseum.org))

## Botany



### Associate Editor

Sabine Huhndorf ([shuhndorf@fieldmuseum.org](mailto:shuhndorf@fieldmuseum.org))

## Geology



### Associate Editor

Olivier Rieppel ([orieppel@fieldmuseum.org](mailto:orieppel@fieldmuseum.org))

## Zoology



### Associate Editor

Janet Voight ([jvoight@fieldmuseum.org](mailto:jvoight@fieldmuseum.org))

**Cover photograph:** Ventral view of skull, young adult *Coryphodon lobatus* (PM39385) from Deardroff Hill *Coryphodon* Quarry, Piceance Creek Basin, Colorado. Photograph by John Weinstein, © 2009 The Field Museum, Image# GEO86493\_13d.



---

---

# FIELDIANA

---

---

## Geology

NEW SERIES, NO. 52

### **A Paleopopulation of *Coryphodon lobatus* (Mammalia: Pantodonta) from Deardorff Hill *Coryphodon* Quarry, Piceance Creek Basin, Colorado**

**Elizabeth M. McGee**

*Department of Biological Sciences  
San Jose State University  
San Jose, CA 95192-0100, U.S.A.  
email: emcgee@email.sjsu.edu*

**William D. Turnbull**

*Department of Geology  
Field Museum  
1400 South Lake Shore Drive  
Chicago, IL 60605-2496, U.S.A.*

Accepted September 25, 2009  
January 8, 2010  
Publication 1554

---

---

PUBLISHED BY FIELD MUSEUM OF NATURAL HISTORY

---

---

Ref  
QE  
51  
.3  
C4f  
n.s. no. 52

1569159

## Table of Contents

ABSTRACT	1
INTRODUCTION	1
ABBREVIATIONS AND INSTITUTIONS	1
DENTAL TERMINOLOGY AND MEASUREMENTS	1
GEOLOGIC SETTING AND HISTORY OF DISCOVERY	1
COMPOSITION AND CONDITION OF THE DEARDORFF HILL <i>CORYPHODON</i> QUARRY ASSEMBLAGE	2
SYSTEMATIC PALEONTOLOGY	4
<i>Coryphodon lobatus</i>	4
AGE DETERMINATION	5
Age Class 1 (young subadult)	6
Age Class 2 (subadult)	7
Age Class 3 (young adult)	7
Age Class 4 (adult)	7
Age Class 5 (advanced/old adult)	8
ASPECTS OF <i>CORYPHODON</i> LIFE HISTORY	8
SOCIALITY IN <i>CORYPHODON</i> AND FUTURE DIRECTIONS	8
ACKNOWLEDGMENTS	9
LITERATURE CITED	9

## List of Illustrations


1. Dental terminology	2
2. Map of Piceance Creek Basin	2
3. Relative representation of postcranial elements	3
4. Metric variation in lower first molar	4
5. Dental eruption sequence for <i>C. lobatus</i>	5
6. (A) <i>Coryphodon lobatus</i> , PM 39436. (B) <i>C. lobatus</i> , PM 39381	6
7. (A) <i>Coryphodon lobatus</i> PM 39605/39665. (B) <i>C. lobatus</i> , PM 39702	6
8. (A) <i>Coryphodon lobatus</i> PM 39385. (B) <i>C. lobatus</i> , PM 35873	7
9. (A) <i>Coryphodon lobatus</i> PM 35870. (B) <i>C. lobatus</i> , PM 35865	7
10. Metric variation in the lower third molar of <i>Coryphodon lobatus</i> from Deardorff Hill <i>Coryphodon</i> Quarry	8

## List of Tables

1. Inventory of postcranial elements	3
2. Summary statistics of molar measurements	5

## Appendices

I. Metric measurements of lower adult dentition of <i>Coryphodon lobatus</i> from Deardorff Hill <i>Coryphodon</i> Quarry	10
II. Metric measurements of upper adult dentition of <i>Coryphodon lobatus</i> from Deardorff Hill <i>Coryphodon</i> Quarry	11
III. Metric measurements of juvenile dentition of <i>Coryphodon lobatus</i> from Deardorff Hill <i>Coryphodon</i> Quarry	12



Digitized by the Internet Archive  
in 2019 with funding from  
Field Museum of Natural History Library

<https://archive.org/details/paleopopulationo52chic>



# A Paleopopulation of *Coryphodon lobatus* (Mammalia: Pantodonta) from Deardorff Hill *Coryphodon* Quarry, Piceance Creek Basin, Colorado

Elizabeth M. McGee and William D. Turnbull

---

## Abstract

A unique early middle Wasatchian paucispecific bone bed from Deardorff Hill in the Piceance Creek Basin of Colorado contains a minimum of 12 individuals of *Coryphodon lobatus* that range in age from subadult “yearlings” to senescent individuals. The preponderance of *Coryphodon* material in this assemblage (92% of the 700+ complete bones represent a single species of *Coryphodon*) argues for a “catastrophic” origin for this assemblage. The Deardorff Hill *Coryphodon* Quarry preserves one of the most complete dental eruption sequences reported to date for *Coryphodon* and allows interpretation of demographic and life history attributes not ordinarily observable, such as evidence of seasonality in births. In addition, females are disproportionate in number to males, further confirming that this species had a polygynous social structure. Mass mortality assemblages are useful in eliciting a better understanding of the range of variation in single populations. Metrically, the molars of *C. lobatus* specimens from Deardorff Hill *Coryphodon* Quarry have coefficients of variation ranging from 4 to 11, which are comparable to metric variation observed in other mass death *Coryphodon* assemblages. An understanding of the range of variation in this highly variable taxon is of particular importance in the taxonomy and phylogenetic relationships of this ubiquitous Eocene mammal.

## Introduction

*Coryphodon* was a large-bodied, subdigitigrade browsing mammal common in North America, Europe, and Asia from the latest Paleocene to the early Eocene. In addition to the ubiquitous occurrence of isolated teeth and partial jaws in fossil outcrops, there are several instances of mass death assemblages of *Coryphodon*, such as Roehler’s *Coryphodon* Catastrophe Quarry (RCCQ) in Wyoming (McGee, 2001, 2002) and AMNH Quarry 242 in New Mexico (Lucas, 1984). Here we report an additional early middle Wasatchian paucispecific assemblage from Deardorff Hill in the Piceance Creek Basin of Colorado. Originally excavated in the 1940s by Bryan Patterson and colleagues, this assemblage provides tantalizing details on intraspecific and interspecific variation in *Coryphodon*. Below we describe the geologic and taphonomic setting of Deardorff Hill *Coryphodon* Quarry and provide a detailed description of the dental material of *Coryphodon lobatus*, including a reconstruction of the eruption sequence of the dentition. As a “death assemblage,” Deardorff Hill *Coryphodon* Quarry provides an unusual opportunity to assess paleobiological and paleodemographic attributes of a population not normally preserved in the fossil record.

## Abbreviations and Institutions

AMNH = American Museum of Natural History, New York, NY; RCCQ = Roehler’s *Coryphodon* Catastrophe Quarry, WY; UCMP = University of California Museum of Paleontology, Berkeley, CA; FMNH or FM = Field Museum, Chicago, IL; PM or P = Field Museum fossil mammal collection.

## Dental Terminology and Measurements

L = length; AW = anterior width; PW = posterior width; MNI = minimum number of individuals; N = number of specimens; CV = coefficient of variation; SD = standard deviation. All measurements are in millimeters. Cusp and loph designations for the M<sub>3</sub> and M<sup>3</sup> are illustrated in Figure 1. Length and width of individual teeth were measured along maximum dimensions in anteroposterior and transverse planes, respectively. Measurements were taken using Fowler Ultra-Cal II electronic calipers.

## Geologic Setting and History of Discovery

Deardorff Hill *Coryphodon* Quarry is situated on the east side of the Piceance Creek Basin in northwestern Colorado. The eastern margin of the Piceance Creek Basin, which formed during the Late Cretaceous, is framed by the Axial Basin arch, White River uplift, and Elk Mountains, while the western margin comprises the Uncompahgre uplift, Douglas Creek arch, and the Uinta arch and was formed during the Late Paleocene and Early Eocene (Figure 2; Kihm, 1984). The three early Tertiary rock units include the Debeque (originally identified as Wasatch by Hayden, 1873), Green River, and the Uinta Formations.

In the 1930s and early 1940s, Bryan “Pat” Patterson of the Field Museum led collecting expeditions to the Plateau Valley region in the Piceance Creek Basin (western Colorado). In 1941, crew member John M. Schmidt discovered Deardorff Hill *Coryphodon* Quarry (“42-41”), which was subsequently excavated by Schmidt and the chief fossil preparator for the Field Museum, James H. Quinn. While Patterson’s work in western Colorado culminated in a number of publications on






FIG. 1. Terminology of the cusps and lochs of the upper and lower molars of *Coryphodon lobatus* as illustrated by a left  $M_3$  and a right  $M_3$ . The terminology used in describing *Coryphodon* lower dentition follows Lucas (1984). Lucas's (1984) terminology is derived from Szalay (1969) and Simpson (1929) with the exception that the cusp identified by Simpson (1929, fig. 8A') as the entoconid is most likely the hypoconulid (Uhen and Gingerich, 1995). This is a significant point since the presence or absence of the entoconid is an important nonmetric trait in distinguishing species of *Coryphodon*. After Lucas (1984, figs. 50F and 51F).

*Titanoides* and other Tertiary mammals, his work on *Coryphodon* was restricted to a single paper (Patterson, 1939) published prior to the discovery of the Deardorff Hill *Coryphodon* Quarry.

Kihm (1984) described the "Plateau Local Fauna" as a series of Middle Clarkforkian (Earliest Eocene) through late Wasatchian (late Early Eocene) faunas. He identified three Tertiary mammal locality sequences in the Piceance Creek Basin: 1) White River in the northern end of the basin and in the Gray Hills region, 2) farther south, the Roan Cliff region (northwest of Battlement Mesa), and 3) the Mamm Creek region (northeast of Battlement Mesa). Deardorff Hill *Coryphodon* Quarry is located in the southeastern margin of the Mamm Creek region. This region, which extends northward to the Grand Hogback mountains and southward to the northeast corner of Battlement Mesa, contains predominantly Clarkforkian to Wasatchian exposures.

### Composition and Condition of the Deardorff Hill *Coryphodon* Quarry Assemblage

The Deardorff Hill *Coryphodon* Quarry is a paucispecific assemblage consisting of over 700 recognizable skeletal elements (not including ribs) and over 600 fragments of *C.*




FIG. 2. Piceance Creek Basin in northwestern Colorado. The extent of the basin is demarcated in light gray. Tertiary mammal locality sequences worked by Kihm (1984) in the Piceance Creek Basin are shown in darker gray and include White River in the northern end of the basin and in the Gray Hills region, the Roan Cliff region (north of Battlement Mesa), and the Mamm Creek region (east of Battlement Mesa). Deardorff Hill *Coryphodon* Quarry is located in the southeastern margin of the Mamm Creek region.

*lobatus*. The MNI based on mandibular and maxillary elements is 10, while MNI based on the right humerus is 12. Information loss is clearly evident in the inventory of bones from this quarry, as depicted in Figure 3 and Table 1, which show the relative abundance of different postcranial elements in the assemblage based on an MNI of 12. The assemblage is biased against smaller elements (e.g., clavicle, sternum, and bones that make up the manus and pes) that, according to the classic taphonomy studies of Voorhies (1969), are the first to be removed by water currents on deposition. Voorhies (1969) suggests that shape of an element influences its potential for transport, so flat bones that float more easily, such as the scapula and innominate, should likewise be absent in an assemblage dominated by larger, heavier bones (i.e., humerus, femur). The profile in Figure 3, however, shows that innominates and scapulae are only slightly less common than larger bones such as the femur. The humerus, the stoutest bone in the *Coryphodon* skeleton, is noticeably more common than any other single element. These biases would suggest that the assemblage was only briefly worked by water currents before deposition and burial.

Fragmentation and compression are characteristic postmortem alterations in the material from Deardorff Hill *Coryphodon* Quarry. Few of the elements were preserved articulated in situ (e.g., PM 39705: atlas through 5th cervical vertebrae; PM 39392: articulated pelvic girdle with left and right innominates, sacrum, and several caudal vertebrae) despite careful preparation of the blocks in the lab at the Field Museum.






FIG. 3. Relative representation of postcranial elements of *Coryphodon lobatus* from Deardorff Hill *Coryphodon* Quarry. Percentages are based on an MNI of 12.

Hill (1979) suggested that the taphonomic profile of an assemblage may reflect the sequence of disarticulation (Hill, 1979). Water, long known to play an important role in the dispersal of bones (Voorhies, 1969; Behrensmeyer, 1975, 1991), also is a factor in determining the disarticulation sequence of a carcass. In humid conditions, for example, a bovid will disarticulate from the extremities inward, whereas in drier conditions, disarticulation proceeds outward from the body to the extremities. If an assemblage is affected by the sequences of bone disarticulation, a bias against proximal or distal bones would be predicted. Counts of distal (i.e., radius, ulna, tibia, and fibula) and proximal (i.e., humerus, scapula, innominate, and femur) elements in the Deardorff Hill *Coryphodon* Quarry were analyzed using the Mann-Whitney test for independence of means. There is no significant difference between the number of proximal vs. distal elements in this assemblage ( $p = 0.686$ ). The assemblage was also analyzed to determine if there were differences between left

and right elements of the limb. Similarly, there was no statistically significant difference between left and right ( $p = 0.931$ ).

Information on relative representation of different elements, the assessment of proximal vs. distal, and left vs. right elements, plus visual inspection of the physical characteristics of the bones, all suggest that burial was rapid and that reworking was minimal.

More than 97% of the assemblage is monospecific. Additional mammals present include primates (*Cantius abditus*: P26477, P26656, P26660), rodents (*Paramys copei*: P26659), creodonts (*Oxyaena forcipata*: P26647), carnivores (*Didymictis protenus*: P26649), condylarths (*Phenacodus primaevus*: P15697; *Hyopsodus* sp. nr. *H. latidens*: P26650, P26697; *Meniscotherium tapiacitis*: P26658), and perissodactyls (*Hyracotherium* sp. C. sensu Kihm 1984: P15705, P26516, P26566, P26567, P26646, P26651, P26655, P26657, P26661, P26664, P26665). The assemblage also contains a few

TABLE 1. Inventory of postcranial elements of *C. lobatus* from Deardorff Hill *Coryphodon* Quarry, and associated relative abundance estimates.

	Number in assemblage	Number per individual	Number expected in assemblage if MNI = 12	Relative abundance
Humerus (L = 11; R = 12)	23	2	24	0.96
Radius (L = 2; R = 3; indet. = 9)	14	2	24	0.58
Scapula (L = 6; R = 4; indet. = 1)	11	2	24	0.46
Ulna (L = 6; R = 7)	13	2	24	0.54
Femur (L = 6; R = 8)	14	2	24	0.58
Innominate (L = 6; R = 4; indet. = 1)	11	2	24	0.46
Tibia (L = 7; R = 5)	12	2	24	0.50
Fibula (L = 0; R = 2; indet. = 3)	5	2	24	0.21
Ribs	259	34	408	0.63
Vertebrae	207	47	564	0.37
Patella (L = 1; R = 3; indet. = 3)	7	2	24	0.29
Manus+pes	225	106	1272	0.18
Clavicle (indet. = 3)	3	2	24	0.13
Sternum	2	6	72	0.03






FIG. 4. Metric variation in the lower first molar of *Coryphodon* from Deardorff Hill *Coryphodon* Quarry compared with lower first molar of *Coryphodon lobatus* reported by Lucas (1984). The Deardorff Quarry specimens slightly extend the size range of *C. lobatus*.

specimens of turtles (*Chelonia*: *Trionychidae*?). The preponderance of *Coryphodon* material in this assemblage (700+ complete bones) and the relative absence of material from other species (21 specimens total, representing eight species of mammals, plus a small assortment of turtle scrap) argues strongly for a “catastrophic” origin for this assemblage.

A catastrophic or mass death accumulation is also characterized by the predominance of a single taxon which shows an age profile that is representative of a living population (Kurtén, 1953; Voorhies, 1969; Turnbull & Martill, 1988). In contrast, assemblages formed over an extended period of time have an age profile that reflects greater mortality in juveniles and old adults (Klein & Cruz-Urbe, 1984; Haynes, 1985, 1987). The Deardorff Hill *Coryphodon* Quarry has an age profile in which each age-group (see below) is represented by about the same number of individuals and is therefore more comparable to a mass death accumulation.

## Systematic Paleontology

Class Mammalia Linnaeus, 1758

Order Pantodonta (Cope, 1873)

Family Coryphodontidae (Marsh, 1876)

*Coryphodon lobatus* (Cope, 1877)

(Figures 1, 5–8)

REFERRED SPECIMENS—P15628: mandible with  $LI_1-M_3$ ,  $RI_1-M_3$  and skull with  $LI^{1-3}$ , LC,  $LP^2-M^3$ ,  $RI^{1-3}$ , RC,  $RP^2-M^3$ ; PM 35865: mandible with  $LI_{1-2}$ ,  $LP_1-M_3$ ,  $RI_2-I_3$ ,  $RP_1-M_3$  and associated lower L/C and  $LI_3$ ; PM 35870: dentary with  $LP_1-M_2$ , associated lower LC and two incisors, and skull with LC,  $LP^{2-4}$ ,  $LM^{1-3}$ ,  $RP^{2-4}$ ,  $RM^{1-3}$ ; PM 35871: skull with  $LI^1-M^3$ ,  $RI^2-C$ ,  $RP^2-M^3$ ; PM 35873: mandible with  $LI_1-M_3$ ,  $RI_1-C$ ,  $RP_2-M_3$ ; PM 35878: mandible with  $LI_1-M_3$ ,  $RI_1-I_2$ , RC- $M_3$ ; PM 35879: maxillary fragment with  $LdP^{2-4}$ ,  $LM^1$ ; PM 35903: skull with  $LP^2-M^3$  and  $RP^2-M^3$ ; PM 39374: mandible with  $LI_{1-2}$ ,  $LP_2-M_3$ ,  $RI_{1-2}$ ,  $RP_1-M_3$ ; PM 39375: mandible

with  $LdP_{1-3}$ ,  $LM_1$  and  $RM_1$ ; PM 39381: maxillary fragment with  $RdP^{1-4}-M^1$  (possibly the other half of PM 39686); PM 39385: skull with LC- $M^3$  and RC,  $RP^2-M^3$ ; PM 39436: mandible with  $LdP_{2-4}-M_1$  (possibly part of PM 35879); PM 39605/39665: skull with  $LI^3?$ ,  $LdP^4$ ,  $LM^{1-2}$  and  $RdP^4$ ,  $RM^{1-2}$ ; PM 39673: skull with  $LI^3$ , LC,  $LP^2-M^3$  and  $RI^3$ , RC,  $RP^2-M^3$ ; PM 39686: maxillary fragment with  $LdP^{1-4}-M^1$ ; PM 39702: mandible with  $LdP_1$ ,  $LdP_{3-4}-M_2$  and  $RI_3?$ ,  $RdP_{1-4}-M_2$ .

DESCRIPTION—*Coryphodon* species have historically been difficult to distinguish (Earle, 1892; Patterson, 1939) because of individual, sexual, and interspecific variation. Lucas’s (1984) treatment remains the most comprehensive to date and provides a basis for identifying the species present at Deardorff Quarry. Lucas (1984) argues that the following characteristics of the  $M^3$  are useful in identifying species of *Coryphodon* (Figure 1): 1) presence and size of  $M^3$  metastyle, 2) relative length and orientation of the postparacrista–premetacrista crest, 3) presence or absence of postprotocrista, and 4) size of the postero-lingual cingulum. For the  $M_3$ , the following characteristics are distinctive (Lucas, 1984): 1) length and orientation of the hypolophid, 2) size of the posterior cingulid, 3) presence or absence of the entoconid, entocristid, and other postero-lingual cusplids and their position and relative size, and 4) size and orientation of the cristid obliqua.

We assign the Deardorff Quarry *Coryphodon* specimens to *C. lobatus*. The  $M_3$  of these specimens possess 1) a semitransverse hypolophid (although not quite as parallel as in *C. subquadratus*), 2) a small but distinct entoconid (absent in *C. molestus*; *C. anthracoides* has numerous lingual cusps), and 3) a well-developed posterior cingulum with an incipient cusp on the lingual side. The hypoconulid in the Deardorff specimens is more lingual (as in other species) than in *C. proterus* in which the hypoconulid is very centrally placed. The  $M^3$  of the specimens from Deardorff Hill *Coryphodon* Quarry 1) typically lack an  $M^3$  metastyle (or otherwise have a very small one), 2) have a distinctive ectoloph because of the orientation of the postparacrista–premetacrista crest, and 3) have a well-developed postprotocrista. Although several of the *Coryphodon* from Deardorff Hill *Coryphodon* Quarry possess a postero-lingual cingulum, Lucas (1984) argues this is not a feature of *C. lobatus*. A large postero-lingual cingulum is seen in *C. proterus*. The Deardorff Hill *Coryphodon* Quarry specimens cannot be *C. proterus*, however, because of the angle formed by the postparacrista–premetacrista; in *C. proterus*, the postparacrista–premetacrista is straight and nearly transverse (Lucas, 1984).

Lucas (1984) assigns Deardorff Hill *Coryphodon* Quarry specimens to two *Coryphodon* species, *C. lobatus* (P15628, PM 35865, PM 35873) and *C. molestus* (PM 35870, PM 35871, PM 35903, and PM 35878). We observe most, if not all, of the characteristics noted above for *C. lobatus* on the specimens Lucas (1984) designates to *C. molestus*. For PM 35870, the upper third molars lack a metastyle, the postprotocrista is present, and there is a distinctive ectoloph. Both PM 35871 and PM 35903 have a very small metastyle on  $M^3$ ; a postprotocrista is present, and the ectoloph is distinctive. PM 35878 has a very small entoconid on the  $M_3$ ; the hypolophid is nearly transverse, and the posterior cingulid is present and well developed. Metrically, PM 35870, PM 35871, PM 35903, and PM 35878 (see below) are smaller than the other Deardorff Hill *Coryphodon* Quarry specimens noted by



TABLE 2. Summary statistics of molar measurements for *C. lobatus* from Deardorff Hill *Coryphodon* Quarry. All measurements in mm.

	N	Mean (mm)	SD	Range (mm)	CV
Lower dentition					
P <sub>4</sub> L	6	23.11	1.58	20.71–24.80	6.84
P <sub>4</sub> W	6	19.40	1.14	17.97–20.58	5.87
M <sub>1</sub> L	9	27.73	2.70	25.08–32.57	9.75
M <sub>1</sub> AW	9	20.45	1.18	18.31–22.66	5.77
M <sub>1</sub> PW	9	20.68	0.88	19.54–22.32	4.27
M <sub>2</sub> L	6	34.84	3.46	30.17–39.47	9.94
M <sub>2</sub> AW	7	25.83	1.59	24.20–28.64	6.16
M <sub>2</sub> PW	6	24.87	1.60	22.97–27.18	6.45
M <sub>3</sub> L	4	38.91	4.46	35.29–44.83	11.45
M <sub>3</sub> AW	5	28.01	1.80	26.24–30.15	6.42
M <sub>3</sub> PW	4	25.25	1.39	23.63–26.94	5.50
Upper dentition					
P <sup>4</sup> L	6	20.15	1.70	18.16–22.58	8.41
P <sup>4</sup> W	6	32.00	1.36	30.64–34.34	4.26
M <sup>1</sup> L	10	27.92	3.03	22.81–31.71	10.84
M <sup>1</sup> AW	10	31.03	1.61	28.77–34.83	5.18
M <sup>1</sup> PW	9	30.14	2.69	26.37–36.15	8.93
M <sup>2</sup> L	7	33.52	2.68	30.28–36.92	7.99
M <sup>2</sup> AW	7	38.40	1.28	37.28–40.54	3.34
M <sup>2</sup> PW	7	37.43	1.70	35.78–40.23	4.53
M <sup>3</sup> L	6	32.74	3.44	28.40–38.40	10.51
M <sup>3</sup> W	6	44.16	3.76	40.14–49.55	8.51

Lucas (1984), but we argue that these specimens should be classified with *C. lobatus*.

*Coryphodon* species have also traditionally been differentiated metrically. An important caveat, however, is that interspecific variation of taxonomic consequence is best assessed using metric measurements on elements that are not sexually dimorphic. Gingerich (1981) suggests that the central cheek teeth (P<sub>4</sub><sup>4</sup>, M<sub>1</sub><sup>1</sup>, and M<sub>2</sub><sup>2</sup>) are unimodally distributed within a species; of the three, the first molar shows the least variation within mammalian species (Gingerich, 1974). A bivariate plot depicting the relationship between length vs. width of the lower first molar for the Deardorff Hill *Coryphodon* Quarry specimens and other *C. lobatus* is presented in Figure 4. The Deardorff Hill *Coryphodon* Quarry assemblage moderately extends the size range for *C. lobatus* with the individuals from Deardorff Hill *Coryphodon* Quarry being smaller on average with respect to other *C. lobatus*.

Coefficients of variation, a standard measure of the relative amount variation in a sample, for P<sub>4</sub> through M<sub>3</sub> of the

*Coryphodon* from Deardorff Hill *Coryphodon* Quarry are given in Table 2. Simpson et al. (1960) noted that coefficients of variation for mammalian species characteristically range between 4 and 10. Gingerich and Schoeninger (1979) observed a more narrow range of approximately 6 to 9 for P<sub>4</sub>, M<sub>1</sub>, M<sub>2</sub>, and M<sub>3</sub> for primates, while Gingerich and Winkler (1979) observed a slightly broader range of approximately 3 to 12 for the same teeth in the red fox (*Vulpes vulpes*). The *Coryphodon* from Deardorff Hill *Coryphodon* Quarry have CV values ranging from approximately 4 to 11. Variation in specimens from Deardorff Hill *Coryphodon* Quarry is within an acceptable range for a species of *Coryphodon*, especially given that this taxon is sexually dimorphic.

### Age Determination

Lucas and Schoch (1990) and Lucas (1984) proposed a dental eruption sequence in *Coryphodon* that is confirmed by the *C. lobatus* specimens from Deardorff Hill *Coryphodon* Quarry. In addition, the specimens from this quarry permit us to fine-tune some of the details of the eruption sequence. We observe the following at Deardorff Hill *Coryphodon* Quarry (Figure 5):

- None of the anterior deciduous dentition is intact in the Deardorff Hill *Coryphodon* Quarry specimens. Lucas and Schoch (1990) believe that the anterior deciduous dentition precedes the posterior deciduous dentition. Given the teeth present in the Deardorff Hill *Coryphodon* Quarry specimens, we believe that the deciduous premolars dP<sub>4</sub>, dP<sub>3</sub>, and dP<sub>2</sub> erupt sequentially, starting with the dP<sub>4</sub>.
- The first molar (M<sub>1</sub>) erupts next at about the same time as I<sub>3</sub> and followed by the last of the deciduous teeth, the dP<sub>1</sub>. Unlike other deciduous teeth, the dP<sub>1</sub> is retained in the adult dentition.
- The M<sub>2</sub> erupts next, followed by the permanent premolars P<sub>4</sub>, P<sub>3</sub>, and P<sub>2</sub>, which replace the deciduous premolars.
- The I<sub>1</sub> probably erupts next, followed by the adult canine, I<sub>2</sub>, and M<sub>3</sub>, which erupt at approximately the same time. From the Deardorff Hill *Coryphodon* Quarry specimens, we can determine that these teeth erupt simultaneously, whereas previously it was suggested that the I<sub>2</sub> and M<sub>3</sub> erupted after the canine.

As in other mammals, deciduous teeth are identified by their weak or flaring roots, a light brown color, and thin, less rugose or striated enamel. Corresponding teeth of the upper and lower dentition erupt at the same time. These patterns and




FIG. 5. Dental eruption sequence for *Coryphodon*. Data from Deardorff Hill *Coryphodon* Quarry confirm postulated sequence of Lucas and Schoch (1990). After Lucas and Schoch (1990).





FIG. 6. (A) *Coryphodon lobatus*, PM 39436, mandible with LdP<sub>2-4</sub>-M<sub>1</sub>. (B) *C. lobatus*, PM 39381, maxillary fragment with RdP<sup>1-4</sup>, M<sup>1</sup>. Deciduous premolars still present and first permanent molar present. I<sub>3</sub> is in the process of erupting (indicated by arrow in Fig. 6A). Deardorff Hill *Coryphodon* Quarry contains four individuals from age class 1 that are nearly identical in eruption and wear patterns. This suggests these individuals were the same age at death and therefore were born at the same time of year (i.e., births could have been seasonal).

characteristics are verified here and also reported by Lucas and Schoch (1990).

We group the *C. lobatus* specimens from Deardorff Hill *Coryphodon* Quarry into five age classes. The first three classes are distinguished by eruption sequences, while the last two classes are differentiated on the basis of wear.

#### Age Class 1 (young subadult)

(Figure 6)

Deciduous premolars are still present, and the first permanent molar is fully erupted; I<sub>3</sub> is in the process of erupting (arrow, Figure 6). Specimens in age class 1 include the following:

PM 39436 (LdP<sub>2-4</sub>-M<sub>1</sub>): M<sub>1</sub> is fully erupted and I<sub>3</sub> is partially erupted. An alveolus is present for dP<sub>1</sub> but the tooth is missing. The dP<sub>2-4</sub> are well worn; the M<sub>1</sub> shows minimum wear, appearing as a shear surface along the metalophid.

PM 39381 (RdCP<sup>1-4</sup>-M<sup>1</sup>): Permanent right I<sub>3</sub> in the crypt exposed through breakage, suggesting that this would be the next tooth to erupt (possibly in advance of M<sub>2</sub>).




FIG. 7. (A) *Coryphodon lobatus*, PM 39605/39665, skull with LI<sup>3?</sup>, LdP<sup>4</sup>, LM<sup>1-2</sup> and RP<sup>3</sup>-M<sup>2</sup>. (B) *C. lobatus*, PM 39702, mandible with LdP<sub>1</sub>, LdP<sub>3-4</sub>-M<sub>2</sub> and RI<sub>3?</sub>, RdP<sub>1-4</sub>-M<sub>2</sub>. These specimens represent age class 2. The second permanent molar is partially erupted; third and fourth permanent premolars in process of erupting (indicated by arrow in Fig. 7A).

PM 35879 (LdP<sub>2-4</sub>-M<sub>1</sub>): Possibly from the same individual as PM 39381. These two specimens are very similar in eruption and wear stage to PM 39436.

PM 39686 (LdP<sup>1-4</sup>-M<sup>1</sup>): Similar in eruption and wear to PM 39436, PM 39381, and PM 35879. This specimen preserves the area posterior to the M<sup>1</sup>, and there is no evidence of the M<sup>2</sup> or even a crypt for M<sup>2</sup>.

PM 39375 (LdP<sub>1-3</sub>, LM<sub>1</sub>, and RM<sub>1</sub>): M<sub>1</sub> and dP<sub>1</sub> are fully erupted. The crypt of the M<sub>2</sub> is visible, but no tooth is apparent. It is also uncertain if the P<sub>4</sub> was shed or fell out postmortem. LdP<sub>1-3</sub> are present with wear (dP<sub>4</sub> is missing, but the alveolus is present and well defined), and there is no wear on M<sub>1</sub>. The presence of the M<sub>2</sub> crypt suggests that this individual was older than PM 39686, PM 39436, PM 39381, and PM 35879.

Deardorff Hill *Coryphodon* Quarry contains four individuals from this class (PM 39686, PM 39436, PM 39381, and PM 35879) that are nearly identical in eruption and wear patterns. This suggests the same age at death and thus would have been born at the same time of year. Although these four are the youngest in the assemblage, the presence of the permanent M<sub>1</sub> suggests that even the youngest individuals in this *Coryphodon* herd were probably weaned and that births could have been seasonal. In two extant analogues of *Coryphodon*—*Sus scrofa* and *Hippopotamus amphibius*—subadults are weaned (3.38 months and 10.13 months, respectively; Ernest, 2003) before the first molar has completely erupted (5.6 months and 24 months, respectively; Smith, 2000).





FIG. 8. (A) *Coryphodon lobatus*, PM 39385, skull with LC-M<sup>3</sup>; RC, P<sup>2</sup>-M<sup>3</sup>. (B) *C. lobatus*, PM 35873, mandible with LI<sub>1</sub>-M<sub>3</sub>, RI<sub>1</sub>-C, P<sub>2</sub>-M<sub>3</sub>. These specimens represent age class 3. Permanent canine and third molar partially erupted, and the adult second incisor (one of the last teeth to erupt in adult dentition) is also erupting.

#### Age Class 2 (subadult)

(Figure 7)

PM 39702 and PM 39605/39665, which may represent the same individual, are considered a subadult:

PM 39702 (mandible with LdP<sub>1</sub>, LdP<sub>3-4</sub>-M<sub>2</sub> and RI<sub>3</sub>, RdP<sub>1-4</sub>-M<sub>2</sub>): M<sub>1</sub> and dP<sub>1-4</sub> are fully erupted and show some wear. The M<sub>2</sub> is partially erupted. On the left, a permanent premolar is partially exposed under the dP<sub>4</sub>. There is a very small alveolus anterior to the right dP<sub>1</sub>, presumably for a deciduous canine; the left side of the mandible is too badly damaged to determine the status of this area. There is a large, spatulate tooth anterior to this alveolus, which is interpreted to be the RI<sub>3</sub>.

PM 39605/39665 (skull with LI<sup>3?</sup>, LdP<sup>4</sup>, LM<sup>1-2</sup>, and RP<sup>3</sup>-M<sup>2</sup>): As with PM 37902, the M<sup>1</sup> shows some wear and the M<sup>2</sup> is partially erupted. Permanent premolars are partially exposed under the LdP<sup>3</sup>, LdP<sup>4</sup>, RdP<sup>3</sup>, and RdP<sup>4</sup>.

#### Age Class 3 (young adult)

(Figure 8)

PM 35873 and PM 39385 are considered young adults:

PM 35873: The C and M<sub>3</sub> are partially erupted on both sides. The anterior dentition of this specimen is unusual in that there are three fully erupted and worn incisors that appear to be part of the adult dentition, but there is also another incisor erupting into the position occupied by RI<sub>2</sub>; it is unknown




FIG. 9. (A) *Coryphodon lobatus*, PM 35870 skull with L/C, LP<sup>2-4</sup>, LM<sup>1-3</sup>, RP<sup>2-4</sup>, RM<sup>1-3</sup>. (B) *C. lobatus*, PM 35865 (mandible with LI<sub>1-2</sub>, P<sub>1</sub>-M<sub>3</sub>, RI<sub>2</sub>-I<sub>3</sub>, P<sub>1</sub>-M<sub>3</sub> and associated lower LC and LI<sub>3</sub> not pictured). All teeth present are part of the permanent dentition (dP<sub>1</sub> is retained). First and second molars are worn (especially the M<sub>1</sub>); incisors also show wear.

whether this tooth is supernumerary or if it is an adult incisor displacing the dI<sub>2</sub>. (This confirms Lucas's observation that the I<sub>2</sub> is the last incisor to erupt.) All other teeth present in the mandible are permanent. The canines are about 28.71 mm (right side) at the gum line, but the dentary is broken away to expose the rest of the canine (on both sides), which appears large. PM 39385 is a skull that similarly has partially erupted M<sup>3</sup> and almost fully erupted canines; the incisors are not preserved in this specimen.

#### Age Class 4 (adult)

(Figure 9)

In age class 4, all of the adult teeth are erupted:

PM 35870 (dentary with LP<sub>1</sub>-M<sub>2</sub>, associated lower LC and two incisors, and skull with LC, LP<sup>2-4</sup>, LM<sup>1-3</sup>, RP<sup>2-4</sup>, RM<sup>1-3</sup>): All teeth are part of the permanent dentition. The protoconids and hypoconids on the lower M<sub>1</sub> and M<sub>2</sub> are worn (especially in M<sub>1</sub>). In the skull, the protocones and postprotocristae of the M<sub>1</sub> and M<sub>2</sub> are similarly worn; the third molar, which is present in the skull but absent in the dentary fragment, shows only the beginning stages of wear. The permanent premolars are not significantly worn in either the skull or the dentary fragment.

PM 35865 (mandible with LI<sub>1-2</sub>, LP<sub>1</sub>-M<sub>3</sub>, RI<sub>2</sub>-I<sub>3</sub>, RP<sub>1</sub>-M<sub>3</sub> and associated lower LC and LI<sub>3</sub>): All teeth are part of the permanent dentition. There is a small amount of wear on the RM<sub>3</sub>; wear increases anteriorly from M<sub>3</sub> to M<sub>1</sub>. P<sub>4</sub> is not worn,



but  $P_3$  is. The left dentary of this mandible is similar in wear features except that the  $P_4$  is more worn than the  $P_3$ , and overall there is considerably more wear on the teeth on the left than on the right. There is a fair amount of wear on  $RI_3$  and to a lesser extent,  $RI_2$ ; on the left side,  $I_1$  shows more wear than  $I_2$ .

PM 35903 (skull with  $LP^2-M^3$  and  $RP^2-M^3$ ): The lower and upper  $M1$  are very worn, and the protoloph and ectoloph on  $M^3$  are fairly worn.

PM 35871 (skull with  $LI^1-M^3$ ,  $RI^2-C$ ,  $RP^2-M^3$ ) and PM 35878 (mandible with  $LI_1-M_3$ ,  $RI_1-I_2$ ,  $RC-M_3$ ): The protocone of the  $M^3$  is moderately worn, along with the protoloph and ectoloph; as with PM 35903, the upper and lower first molar are very worn. The left and right  $I^3$  in PM 35871 are the most worn of the upper incisors; on the lower dentition (PM 35878), the  $I_2$  are the most worn of the incisors.

#### Age Class 5 (advanced/old adult)

As with age class 4, in age class 5, all of the adult teeth are erupted. Age class 5 exhibits extreme wear on many of the teeth:

PM 39673 (skull with  $LI^3$ ,  $LC$ ,  $LP^2-M^3$  and  $RI^3$ ,  $RC$ ,  $RP^2-M^3$ ), PM 39374 ( $LI_{1-2}$ ,  $LP_2-M_3$ , and  $RI_{1-2}$ ,  $RP_1-M_3$ ), and P15628 (mandible with  $LI_1-M_3$ , and  $RI_1-M_3$  and skull with  $LI^{1-3}$ ,  $LC$ ,  $LP^2-M^3$ ,  $RI^{1-3}$ ,  $RC$ ,  $RP^2-M^3$ ) are the most worn of the specimens. Both upper and lower  $M1$  and  $M2$  are extremely worn, and  $M3$  is beginning to show excessive wear along the metalophid and protoconid/hypolophid on the lowers; in the uppers, the protoloph and ectoloph are completely worn. There is still enamel present elsewhere on the occlusal surface of both the upper and lower  $M3$ ; there is almost no enamel left on the occlusal surfaces of the upper and lower  $M1$  and  $M2$ . The  $P3$  and  $P4$  are also well worn (especially in the lower dentition).

#### Aspects of *Coryphodon* Life History

The preservation of the eruption sequence of the Deardorff Hill *Coryphodon* provides an opportunity to consider life history characteristics of *Coryphodon*. Smith (2000) suggests that relative eruption sequences of select extant taxa can be used to predict the tempo of life histories for species of extinct mammals. She examined eruption sequences in Insectivora, Archonta, and Ungulata to test "Schultz's Rule" that permanent incisors, canines, and premolars (= replacement teeth) erupt earlier in slow-growing, long-lived species. She noted that in rapidly growing mammals (e.g., *Antidorcas*), the three sets of teeth (deciduous, molars, and replacement) erupt sequentially (deciduous → molars → replacement), whereas in slow-growing mammals (e.g., *Homo*), the eruption of molars and replacement teeth is mixed (e.g., the permanent canine erupts after  $M1$  and before  $M3$ , as in the case of the collared peccary, *Tayassu tajacu*). Age classes 1 and 2 of *C. lobatus* from Deardorff Hill *Coryphodon* Quarry clearly indicate a sequence in which the molars and replacement teeth are mixed, a pattern that is suggestive of a prolonged development.

Framing *Coryphodon* evolution within the context of life histories, as suggested by Smith (2000) also leads us to consider which, if any, large-bodied extant herbivore is a good analogue for this taxon. Within the "slow growers," Smith (2000)




FIG. 10. Metric variation in the lower third molar of *Coryphodon lobatus* from Deardorff Hill *Coryphodon* Quarry compared with lower third molar of *C. anthracoides* from Roehler's *Coryphodon* Catastrophe Quarry. Both death assemblages indicate the presence of large- and small-size morphs in each assemblage, presumed to indicate males and females, respectively.

suggested that *Coryphodon* is less like the hippo and more like a large deer or pig. She noted that the eruption patterns of  $I_1$  and  $P_3$  seen in *Coryphodon* are found in extant mammals in which the  $M_1$  erupts between 0.34 and 0.88 years and that have a life span of 21 to 35 years. She questions whether the hippo, with a slower life history marked by the emergence of the  $M1$  at two years and a life span of 40 to 50 years, is a suitable analogue for *Coryphodon*. *Coryphodon* shares a sequence pattern  $M_1M_2P_3M_3$  with *Equus burchelli*, *T. tajacu*, *S. scrofa*, *H. amphibius*, *Ceratotherium simum*, and *Procavia capensis*. We add to this the observation that all these taxa, with the exception of *E. burchelli*, wean their young before the complete eruption of the first molar (Smith, 2000; Ernest, 2003). These observations serve to underscore the complexity of identifying an analogue. In putting *Coryphodon* in the context of an extant species, Lucas (1984) discussed functional analogues such as the pygmy hippo (*Hexaprotodon liberiensis*), tapirs (*Tapirus* sp.), and the Sumatran rhino (*Dicerorhinus sumatrensis*) as well as ecologic analogues such as the hippo (*H. amphibius*). Dental eruption data suggest that life history can also serve as a basis for analogy between extant and extinct taxa.

#### Sociality in *Coryphodon* and Future Directions

Death or catastrophic assemblages such as Deardorff Hill *Coryphodon* Quarry potentially provide a unique window into the behavior of extinct populations. A paleopopulation, defined as a well-delimited faunal level spanning a short period of geological time (MacFadden, 2008), may still represent as much as 100,000 years in duration. Mass death assemblages, representing near instantaneous accumulation (e.g., such as that seen over a period of days, weeks, or months during wildebeest migrations when rivers are in flood; Talbot & Talbot 1963), are more comparable to extant populations.

Mihlbachler (2003) argued that because adult sex ratios (ASRs) are linked to sociality in extant species (Berger, 1986;



Byers, 1997), ASRs in fossil assemblages can be used to infer paleodemography in extinct species if males can be distinguished from females through sexually dimorphic features. Sexual dimorphism in *Coryphodon* quarry assemblages is evident in the size of the third molars and the canines (Figure 10). We would expect this to be the case on the basis of Gingerich's (1974) suggestion that the third molar is under greater influence of sex hormones than the first and second molars, which erupt earlier. Sexual dimorphism can also be inferred from canines. However, canines are often missing or broken in *Coryphodon* assemblages that have undergone significant postmortem modifications. Figure 10 indicates that there are two distinct size morphs in the Deardorff Hill *Coryphodon* Quarry assemblage, a pattern that is also seen in the death assemblages of *C. anthracoides* from Roehler's *Coryphodon* Catastrophe Quarry and *C. molestus* from AMNH Quarry 242. General consensus is that in extant large herbivorous mammals, males are usually larger than females (Ralls, 1977; Jarman, 1983), and this has been inferred for fossil mammals (Kurtén, 1969; Gingerich, 1981).

Ralls (1977) and Jarman (1983) argued that there is a close correlation between sexual dimorphism and polygyny. Using information on sexual dimorphism and body size differences in *C. lobatus* from Deardorff Hill *Coryphodon* Quarry, we can reconstruct a picture of sociality patterns. The ASR at Deardorff Hill *Coryphodon* Quarry is minimally 1:3 (which is also the case for *C. anthracoides* at Roehler's *Coryphodon* Catastrophe Quarry). We would predict based on the model of Berger et al. (2001) that *C. lobatus* lived in unisex groups except during periods of the year when mating took place. The ASR of the Deardorff Hill *Coryphodon* Quarry indicates that males were present at the time the death event occurred, and therefore this assemblage probably preserves a picture of this population of *Coryphodon* during mating season.

The cranial and postcranial *C. lobatus* material from Deardorff Hill *Coryphodon* Quarry is still largely unstudied beyond basic inventorying, and thus many questions remain regarding the degree and nature of sexual dimorphism in this species. It has been suggested that extreme polygyny and dimorphism are associated with bimaturism; that is, males and females reach sexual maturity at different ages (Ralls, 1977; Jarman, 1983). Jarman (1983) argued that a delay in male maturation is in fact a requirement for the evolution of extreme polygyny and sexual dimorphism. *Coryphodon* mass death assemblages will no doubt be useful in eliciting the degree of sexual dimorphism in this taxon and hence the possibility that males matured later than females. Additional work might also focus on a closer inspection of size differences between the two morphs (i.e., males and females) in the Deardorff Hill *Coryphodon* Quarry assemblage. With respect to interpreting the behavioral and/or demographic significance of sexual dimorphism, the distinction must be made between dimorphism in secondary sexual characteristics and dimorphism in body size. A number of authors have noted that selection pressures including sexual selection may contribute to sexual dimorphism in size (Ralls, 1977; Alexander et al., 1979; Gingerich, 1981; Jarman, 1983; Weckerly, 1998). Ralls (1977) suggests that sexual dimorphism in structures and coloration is more closely related to sexual selection than is body size. That is, different factors can cause dimorphism in size (e.g., neonate size in females), but only sexual selection can affect structure/coloration. Future analysis might test whether *Coryphodon* males and females are different only in

size or if differences in morphology beyond allometry exist. If males are relatively and absolutely larger than females, then it may be the case that sexual selection is the driving force.

## Acknowledgments

We would like to thank Bill Simpson (Field Museum) for his invaluable assistance with this project, and John Weinstein (Field Museum) for excellent photographic support. Drs. Spencer Lucas (New Mexico Museum of Natural History and Science) and Lawrence Flynn (Peabody Museum of Archaeology and Ethnology) for excellent feedback on the manuscript. Many thanks to Dr. Allen Kihm for advice on the map of the Piceance Creek Basin and to Arnaud Bergerol for assistance with data entry.

## Literature Cited

- ALEXANDER, R. D., J. L. HOOGLAND, R. D. HOWARD, K. M. NOONAN, AND P. W. SHERMAN. 1979. Sexual dimorphism and breeding systems in pinnipeds, ungulates, primates, and humans, pp. 402–435. In Chagnon, N. A., and W. Irons, eds., *Evolutionary Biology and Human Social Behavior*. Duxbury Press, North Scituate, Massachusetts.
- BEHRENSMEYER, A. K. 1975. The taphonomy and paleoecology of Plio-Pleistocene vertebrate assemblages east of Lake Rudolf, Kenya. *Bulletin of the Museum of Comparative Zoology*, **146**: 473–578.
- . 1991. Terrestrial vertebrate accumulations, pp. 291–335. In Alison, P. A., and D. E. G. Briggs, eds., *Releasing the Data Locked in the Fossil Record*. Plenum Press, New York.
- BERGER, J. 1986. *Wild Horses of the Great Basin: Social Competition and Population Size*. University of Chicago Press, Chicago.
- BERGER, J., S. DULAMTSEREN, S. CAIN, D. ENKKHBILEG, P. LICHTMAN, Z. NAMSHIR, G. WINGARD, AND R. READING. 2001. Back-casting sociality in extinct species: New perspectives using mass death assemblages and sex ratios. *Proceedings of the Royal Society of London B*, **268**: 131–139.
- BYERS, J. A. 1997. *American Pronghorn: Social Adaptations and the Ghosts of Predators Past*. University of Chicago Press, Chicago.
- COPE, E. D. 1873. On the short-footed Ungulata of the Eocene of Wyoming. *Proceedings of the American Philosophical Society*, **13**: 38–74.
- . 1877. Report upon the extinct Vertebrata obtained in New Mexico by parties of the expedition of 1874. Report to the U.S. Geographical Survey West of the 100th Meridian (Wheeler) **4**, **2**: 1–370.
- EARLE, C. 1892. Revision of the species of *Coryphodon*. *Bulletin of the American Museum of Natural History*, **4**: 149–166.
- ERNEST, S. K. M. 2003. Life history characteristics of placental nonvolant mammals. *Ecology*, **84**: 3402.
- GINGERICH, P. D. 1974. Size variability of the teeth in living mammals and the diagnosis of closely related sympatric fossil species. *Journal of Paleontology*, **48**: 895–903.
- . 1981. Variation, sexual dimorphism, and social structure in the early Eocene horse *Hyracotherium* (Mammalia, Perissodactyla). *Paleobiology*, **7**: 443–455.
- GINGERICH, P. D., AND M. J. SCHOENINGER. 1979. Patterns of tooth size variability in the dentition of primates. *American Journal of Physical Anthropology*, **81**: 457–466.
- GINGERICH, P. D., AND D. A. WINKLER. 1979. Patterns of variation and correlation in the dentition of the red fox, *Vulpes vulpes*. *Journal of Mammalogy*, **60**: 691–704.
- HAYDEN, F. V. 1873. Third annual report of the U.S. Geological Survey of the Territories embracing Colorado and New Mexico, for the year 1869, 261 pp.



- HAYNES, G. 1985. Age profiles in elephant and mammoth bone assemblages. *Quaternary Research*, **24**: 333–345.
- . 1987. Proboscidean die-offs and die-outs: Age profiles in fossil collections. *Journal of Archaeological Science*, **14**: 659–668.
- HILL, A. 1979. Disarticulation and scattering of mammal skeletons. *Paleobiology*, **5**: 261–274.
- JARMAN, P. 1983. Mating system and sexual dimorphism in large, terrestrial, mammalian herbivores. *Biological Reviews*, **58**: 485–520.
- KIHM, A. J. 1984. Early Eocene Mammalian Faunas of the Piceance Creek Basin Northwestern Colorado. Ph.D. Thesis, University of Colorado, Boulder.
- KLEIN, R. G., AND K. CRUZ-URIBE. 1984. *The Analysis of Animal Bones from Archeological Sites*. University of Chicago Press, Chicago.
- KURTÉN, B. 1953. On the variation and population dynamics of fossil and recent mammal populations. *Acta Zoologica Fennica*, **76**: 1–122.
- . 1969. Sexual dimorphism in fossil mammals, pp. 226–233. In Westermann, G. E. G., ed., *Sexual Dimorphism in Fossil Metazoa and Taxonomic Implications*. International Union of Geological Sciences, Series A. Schweizerbart'sche Verlagbuchhandlung, Stuttgart, West Germany.
- LUCAS, S. G. 1984. Systematics, Biostratigraphy, and Evolution of Early Cenozoic *Coryphodon* (Mammalia, Pantodonta). Ph.D. Thesis, Yale University, New Haven, Connecticut.
- LUCAS, S. G., AND R. M. SCHOCH. 1990. Ontogenetic studies of early Cenozoic *Coryphodon* (Mammalia, Pantodonta). *Journal of Paleontology*, **64**: 831–841.
- MACFADDEN, B. J. 2008. Geographic variation in diets of ancient populations of 5-million-year-old (early Pliocene) horses from southern North America. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **266**: 83–94.
- MARSH, O. C. 1876. On some characters of the genus *Coryphodon*. *American Journal of Science*, **3**: 425–428.
- MCGEE, E. M. 2001. A mass death accumulation of *Coryphodon anthracoideus* (Pantodonta: Mammalia) from Roehler's *Coryphodon* Catastrophe Quarry (Lower Eocene, Wasatch Formation, Washakie Basin, Wyoming), pp. 317–333. In Gunnell, G. G., ed., *Eocene Biodiversity: Unusual Occurrences and Rarely Sampled Habitats*. Kluwer Academic/Plenum Publishers, New York.
- . 2002. Intraspecific dental variability in cf. *Coryphodon anthracoideus* (Mammalia: Pantodonta) from Roehler's *Coryphodon* Catastrophe Quarry, Washakie Basin, Wyoming. *Rocky Mountain Geology*, **37**: 61–73.
- MIHLBACHLER, M. C. 2003. Demography of late Miocene rhinoceroses (*Teleoceras proterum* and *Aphelops malacorhinus*) from Florida: Linking mortality and sociality in fossil assemblages. *Paleobiology*, **29**: 412–428.
- PATTERSON, B. 1939. A skeleton of *Coryphodon*. *Proceedings of the New England Zoological Club*, **17**: 97–110.
- RALLS, K. 1977. Sexual dimorphism in mammals: Avian models and unanswered questions. *The American Naturalist*, **111**: 917–938.
- SIMPSON, G. G. 1929. A new Paleocene Uintathere and molar evolution in the Amblypoda. *American Museum Novitates*, **387**: 1–9.
- SIMPSON, G. G., A. ROE, AND R. C. LEWONTIN. 1960. *Quantitative Zoology*. Harcourt, Brace, and World, New York.
- SMITH, B. H. 2000. "Schultz's Rule" and the evolution of tooth emergence and replacement patterns in primates and ungulates, pp. 212–227. In Ferguson, M. W. J., ed., *Development, Function and Evolution of Teeth*. Cambridge University Press, New York.
- SZALAY, F. S. 1969. Mixodectidae, Microsypidae, and the insectivore-primate transition. *Bulletin of the American Museum of Natural History*, **140**: 193–330.
- TALBOT, L. M., AND M. H. TALBOT. 1963. The wildebeest in Western Masailand, East Africa. *Wildlife Monographs*, **12**: 5–88.
- TURNBULL, W. D., AND D. M. MARTILL. 1988. Taphonomy and preservation of a monospecific titanotheres assemblage from the Washakie Formation (Late Eocene), southern Wyoming: An ecological accident in the fossil record. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **63**: 91–108.
- UHEN, M. D., AND P. D. GINGERICH. 1995. Evolution of *Coryphodon* (Mammalia, Pantodonta) in the late Paleocene and early Eocene of northwestern Wyoming. *Contributions from the Museum of Paleontology, University of Michigan*, **29**: 259–289.
- VOORHIES, M. 1969. Taphonomy and population dynamics of an early Pliocene vertebrate fauna, Knox County, Nebraska. *Contributions to Geology, University of Wyoming Special Paper*, **1**: 1–69.
- WECKERLY, F. W. 1998. Sexual size dimorphism: Influence of mass and mating systems in the most dimorphic mammals. *Journal of Mammalogy*, **79**: 33–52.

### Appendix I. Metric measurements of lower adult dentition of *Coryphodon lobatus* from Deardorff Hill *Coryphodon* Quarry (all measurements in mm)

	PM 35878		P15628		PM 35865		PM 35873		PM 35870	PM 39374	
	Left	Right	Left	Right	Left	Right	Left	Right	Left	Left	Right
I <sub>1</sub>	18.51		21.79	20.73	20.02		22.51	21.04		20.59	17.51
I <sub>2</sub>	21.45	21.53	21.85	23.68	22.92	23.45	23.02	22.94		22.68	23.09
I <sub>3</sub>	20.51		19.54	18.23	23.26	19.48	24.90	23.95			
C-bl	21.52	22.21	27.66	28.03	26.32						
C-md	24.62	25.80	34.53	33.12	23.97						
p <sub>1</sub> L	15.93	15.99	16.21	16.39	16.95	16.04	17.13		15.86		15.57
p <sub>1</sub> W	9.51	11.06	10.47	11.66	9.73	9.61	9.37		9.45		9.45
p <sub>2</sub> L	17.05	18.59	24.92	20.83	19.34	20.35	22.16	21.27	17.47	18.80	21.41
p <sub>2</sub> W	13.50	13.25	15.62	16.42	13.97	14.17	15.08	14.78	13.41	13.79	13.59
p <sub>3</sub> L	21.43	20.50	23.80	23.50	20.78	20.35	22.60	22.15	22.97	22.34	22.07
p <sub>3</sub> W	15.72	17.39	17.89	17.49	16.31	16.24	17.73	17.63	15.52	16.22	16.10
p <sub>4</sub> L	21.14	18.24	24.51	25.24	23.68	24.13	23.20	23.82	21.73	23.71	23.51
p <sub>4</sub> W	18.49	17.82	20.03	20.55	18.33	19.18	20.44	19.54	17.97	20.27	19.33
m <sub>1</sub> L	25.72	24.34	30.69	29.56	25.29	28.69	29.01	29.99	25.08	25.08	24.71
m <sub>1</sub> AW	20.58	19.06	20.47	20.89	18.92	19.40	20.72	19.64	20.49	22.80	21.70
m <sub>1</sub> PW	21.49	19.68	21.44	25.53	20.42	20.16	21.12	20.44	19.96	21.16	20.61
m <sub>2</sub> L	35.24	31.76	36.09	37.12	35.70	37.18	39.18	38.60	30.17	32.42	29.60
m <sub>2</sub> AW	23.78	24.92	26.26	31.65	24.73	24.43	25.99	25.61	24.20	27.08	24.58
m <sub>2</sub> PW	24.58	24.91	26.67	28.57	23.48	24.61	26.47	26.17	22.97	25.05	22.95
m <sub>3</sub> L	34.87	37.68	45.93	42.32	39.84	39.48				36.49	35.62
m <sub>3</sub> AW	26.82	27.44	31.01	27.42	26.00	26.21	29.84	26.48		27.28	27.09
m <sub>3</sub> PW	25.54	25.04	26.45	28.02	23.55	23.49				24.87	24.14



**Appendix II. Metric measurements of upper adult dentition of *Coryphodon lobatus* from Deardorff Hill  
Coryphodon Quarry (all measurements in mm)**

	PM 35871		PM 39385		PM 35903		PM 39673		PM 35870		P15628	
	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right
I1	21.81										24.54	24.97
I2	21.31	20.06									25.90	23.27
I3	19.63	17.88					19.01	15.22			23.35	22.35
C-bl	23.42	23.57	27.40	27.27			23.46	21.86	29.75		31.13	32.95
C-md	21.93	21.56	19.48	18.69			21.27	21.68	23.81		29.02	30.54
p <sup>1</sup> L	14.31											
p <sup>1</sup> W	10.80		11.20									
p <sup>2</sup> L	18.14	18.37	19.34	19.22	21.82	21.85	20.18	19.85	18.05	18.61	20.53	20.39
p <sup>2</sup> W	28.56	26.64	28.74	29.21	24.80	26.49	27.64	28.50	27.11	26.40	28.59	26.79
p <sup>3</sup> L	19.78	18.87	21.75	21.56	21.81	20.88	19.34	19.63	19.33	19.63	23.49	23.64
p <sup>3</sup> W	30.21	30.61	30.99	31.43	28.70	29.15	31.41	29.33	29.71	29.09	33.24	32.52
p <sup>4</sup> L	18.47	17.83	20.98	20.43	19.77	18.90	17.99	18.46	20.46	20.55	22.51	22.42
p <sup>4</sup> W	31.39	31.45	32.50	31.66	30.96	30.65	30.59	31.70	31.41	32.15	34.58	33.32
m <sup>1</sup> L	22.65	22.90	30.39	28.43	29.09	27.19	23.23			27.03	32.03	30.78
m <sup>1</sup> AW	30.92	30.11	30.54	30.07	31.03	29.08	32.19		27.79	31.43	34.83	
m <sup>1</sup> PW	29.58	29.48	30.79	30.09	29.64	27.93	30.56		30.23	31.49	37.32	33.17
m <sup>2</sup> L	30.49	30.81	37.75	35.38	34.09	35.14	34.66		31.56	31.60	36.58	34.72
m <sup>2</sup> AW	37.22	38.11	37.33	37.99	38.41	37.64	40.54		37.70	37.65	39.75	38.89
m <sup>2</sup> PW	35.43	35.05	34.97	35.54	36.38	35.69	38.26		35.76	35.69	39.87	39.68
m <sup>3</sup> L	27.72	28.43	32.89	31.45	29.46	31.38	33.41	32.25	32.63	33.40	39.47	35.47
m <sup>3</sup> W	40.58	40.22	43.74	44.65	40.54	42.46	45.27	44.43	39.73	43.70	49.55	46.26

**Appendix III. Metric measurements of juvenile dentition of *Coryphodon lobatus* from Deardorff Hill  
Coryphodon Quarry (all measurements in mm)**

	PM 39375 left	PM 39375 right	PM 39702 left	PM 39702 right	PM 39436 left
I <sub>1</sub>					
I <sub>2</sub>					
I <sub>3</sub>				20.30	
C-bl					
C-md					
dp <sub>1</sub> L	16.03		14.15	15.46	
dp <sub>1</sub> W	9.80		9.83	10.75	
dp <sub>2</sub> L	12.16				15.78
dp <sub>2</sub> W	8.56				10.09
dp <sub>3</sub> L	17.60		18.15		19.11
dp <sub>3</sub> W	10.77		11.94		11.52
dp <sub>4</sub> L			20.81		26.71
dp <sub>4</sub> AW			14.79		14.17
dp <sub>4</sub> PW			17.30		15.69
m <sub>1</sub> L	29.75	29.99	25.68	25.33	32.57
m <sub>1</sub> AW	18.64	17.77	21.26	19.13	20.61
m <sub>1</sub> PW	19.48	19.54	19.59	19.35	19.91
m <sub>2</sub> L					
m <sub>2</sub> AW			26.07	23.76	
m <sub>2</sub> PW					
m <sub>3</sub> L					
m <sub>3</sub> AW					
m <sub>3</sub> PW					

	PM 39605/39665 left	PM 39605/39665 right	PM 39686 left	PM 39381 right	PM 35879 left
I <sup>1</sup>					
I <sup>2</sup>					
I <sup>3</sup>	20.42				
C-bl					
C-md					
dp <sup>1</sup> L			16.00	16.34	
dp <sup>1</sup> W			13.91	10.54	
dp <sup>2</sup> L			14.98	17.71	14.67
dp <sup>2</sup> W				15.32	14.55
dp <sup>3</sup> L					16.95
dp <sup>3</sup> W			18.87	18.26	17.01
dp <sup>4</sup> L		22.92	20.69	23.31	20.82
dp <sup>4</sup> W		25.64	24.79	23.73	23.29
m <sup>1</sup> L	27.42	27.56	27.69	29.29	31.21
m <sup>1</sup> AW	30.74	31.05	30.44	30.16	30.33
m <sup>1</sup> PW	28.71	31.15		26.37	30.54
m <sup>2</sup> L	30.33	31.31			
m <sup>2</sup> AW	36.47	37.00			
m <sup>2</sup> PW	38.27	35.32			
m <sup>3</sup> L					
m <sup>3</sup> W					







Field Museum of Natural History  
1400 South Lake Shore Drive  
Chicago, Illinois 60605-2496  
Telephone: (312) 665-7769